Mon. Not. R. Astron. Soc. **000**, 1–14 (2010) Printed 21 December 2011 (MN LATEX style file v2.2)

Photon Geodesics in FRW Cosmologies

Ojeh Bikwa,[1] Fulvio Melia[2⋆] and Andrew Shevchuk[3]†

[1]*Amherst College, Amherst MA 01002, USA*

[2]*Department of Physics, The Applied Math Program, and Department of Astronomy, The University of Arizona, Tucson AZ 85721, USA*

[3]*Department of Astronomy, The University of Arizona, Tucson AZ 85721, USA*

arXiv:1112.4774v1 [astro-ph.CO] 20 Dec 2011

ABSTRACT

The Hubble radius is a particular manifestation of the Universe's gravitational horizon, $R_{\rm h}(t_0) \equiv c/H_0$, the distance beyond which physical processes remain unobservable to us at the present epoch. Based on recent observations of the cosmic microwave background (CMB) with WMAP, and ground-based and HST searches for Type Ia supernovae, we now know that $R_{\rm h}(t_0)$ ~13.5 Glyr. This coincides with the maximum distance ($ct_0 \approx 13.7$ Glyr) light *could* have traveled since the big bang. However, the physical meaning of $R_{\rm h}$ is still not universally understood or accepted, though the minimalist view holds that it is merely the proper distance at which the rate of cosmic recession reaches the speed of light c. Even so, it is sometimes argued that we can see light from sources beyond $R_{\rm h}$, the claim being that $R_{\rm h}$ lies at a redshift of only ~2, whereas the CMB was produced at a much greater redshift (~1100). In this paper, we build on recent developments with the gravitational radius by actually calculating null geodesics for a broad range of FRW cosmologies, to show—at least in the specific cases we consider here, including ΛCDM—that no photon trajectories reaching us today could have ever crossed $R_{\rm h}(t_0)$. We therefore confirm that the current Hubble radius, contrary to a commonly held misconception, is indeed the limit to our observability. We find that the size of the visible universe in ΛCDM, measured as a proper distance, is approximately $0.45ct_0$.

Key words: cosmic microwave background, cosmological parameters, cosmology: observations, cosmology: redshift, cosmology: theory, distance scale

⋆ John Woodruff Simpson Fellow. E-mail: fmelia@email.arizona.edu

† NSF Graduate Fellow.

1 INTRODUCTION

The standard model of cosmology is confronted with several unpalatable coincidences, suggesting that we do not yet have a fully consistent picture of the Universe's dynamical expansion (see, e.g., Melia & Shevchuk 2011). Part of the problem is that cosmological observations can only be interpreted from within the context of a pre-assumed model, and the data can be quite compliant.

The ΛCDM (Cold Dark Matter with a cosmological constant Λ) model has been without peer in cosmology (see, e.g., Spergel et al. 2003, and Tegmark et al. 2004). For example, this model has been used with complementary measurements of the cosmic microwave background (CMB) radiation to determine that the Universe is flat, so its energy density ρ is at (or very near) the "critical" density $\rho_{\rm c} \equiv 3c^2H^2/8\pi G$. But among the many peculiarities of this description of the universe is the inference, based on current observations, that the density $\rho_{\rm d}$ of dark energy must itself be of order $\rho_{\rm c}$. (In the context of ΛCDM, the best fit to the WMAP data indicates that dark energy represents approximately 73% of the total $\rho \approx \rho_{\rm c}$; see Spergel et al. 2003.) Dark energy is often thought to be the manifestation of the aforementioned cosmological constant, Λ, though no reasonable explanation has yet been offered as to why such a fixed, universal density ought to exist at this scale. It is well known that if Λ is associated with the energy of the vacuum in quantum theory, it should have a scale representative of phase transitions in the early Universe—120 orders of magnitude greater than $\rho_{\rm c}$.

Many workers have attempted to circumvent these difficulties by proposing alternative forms of dark energy, including Quintessence (Ratra & Peebles 1988; Wetterich 1988), which represents an evolving canonical scalar field with an inflation-inducing potential, a Chameleon field (see, e.g., Mota & Barrow 2004; Khoury & Weltman 2004; Brax et al. 2004) in which the scalar field couples to the baryon energy density and varies from solar system to cosmological scales, and modified gravity, arising out of both string motivated, or General Relativity modified actions (Capozziello et al. 2003; Nojiri & Odintsov 2003; Carroll et al. 2004), which introduce large length scale corrections modifying the late time evolution of the Universe. The actual number of suggested remedies is far greater than this small, illustrative sample.

An equally perplexing puzzle with ΛCDM has been dubbed the "coincidence problem," arising from the peculiar near-simultaneous convergence of the matter energy density $\rho_{\rm m}$ and $\rho_{\rm d}$ towards $\rho_{\rm c}$ in the present epoch. Though $\rho_{\rm m}$ and $\rho_{\rm d}$ are expected to change at different rates as the Universe expands (particularly if dark energy is a cosmological constant) they are nearly equal in the present epoch, implying that we live at a special time in cosmic history.

In a recent paper (Melia & Shevchuk 2011), we proposed an explanation for yet another disturbing coincidence, having to do with the apparent equality of our gravitational (or Hubble) radius $R_{\rm h}$ with the distance ct_0 light could have traveled since the big bang (in terms of the presumed current age t_0 of the Universe). This equality has received some scrutiny in recent years (Melia 2003, 2007, 2009, Melia & Abdelqader 2009, van Oirschot et al. 2010; see also Lima 2007 for a related, though unpublished, work).

Unfortunately, there is still some confusion regarding the properties of $R_{\rm h}$ due to a misunderstanding of the role it plays in our observations. For example, it is sometimes suggested (see, e.g., Davis & Lineweaver 2004; van Oirschot et al. 2010) that sources beyond $R_{\rm h}(t_0)$ are observable today, which is certainly not the case. We will therefore begin by elaborating upon what the gravitational radius $R_{\rm h}$ is—and what it is not. Though first defined in Melia (2007), an unrecognized form of $R_{\rm h}$ actually appeared in de Sitter's (1917) own account of his spacetime metric. And we will advance the discussion further by actually calculating photon trajectories for various well-studied Friedmann-Robertson-Walker (FRW) Cosmologies, demonstrating that the null geodesics reaching us at t_0 have never crossed $R_{\rm h}(t_0)$. Some come close, and in one case—the de Sitter model—they approach $R_{\rm h}$ asymptotically as t recedes to our infinite past. Our conclusion in this paper will be that $R_{\rm h}(t_0)$ is a real limit to our observability at the present time t_0.

2 THE GRAVITATIONAL (HUBBLE) RADIUS

Standard cosmology is based on the Robertson-Walker metric for a spatially homogeneous and isotropic three-dimensional space. In terms of the proper time t measured by a comoving observer, and the corresponding radial (r) and angular coordinates (θ and ϕ) in the comoving frame, an interval ds in this metric is written as

$$ds^2 = c^2\,dt^2 - a^2(t)[dr^2(1-kr^2)^{-1} + r^2(d\theta^2 + \sin^2\theta\,d\phi^2)]\ , \qquad (1)$$

where $a(t)$ is the expansion factor and the constant k is $+1$ for a closed universe, 0 for a flat, open universe, or -1 for an open universe.

In recent work (Melia 2007, 2009; Melia & Abdelqader 2009), we demonstrated the usefulness of examining properties of the metric in terms of both co-moving coordinates (ct, r, θ, ϕ) and observer-dependent coordinates (cT, R, θ, ϕ), where R is the so-called proper radius $a(t)r$. Whereas (ct, r, θ, ϕ) describe events in a frame "falling" freely with the cosmic expansion, the second set

of coordinates are referenced to a particular individual who describes the spacetime relative to the origin at his location.[1]

For a flat universe ($k = 0$), it is straightforward to show that with this definition of R, Equation (1) becomes

$$ds^2 = \Phi\left[c\,dt + \left(\frac{R}{R_{\rm h}}\right)\Phi^{-1}dR\right]^2 - \Phi^{-1}dR^2 - R^2\,d\Omega^2\ , \tag{2}$$

where the function

$$\Phi \equiv 1 - \left(\frac{R}{R_{\rm h}}\right)^2 \tag{3}$$

signals the dependence of the metric on the proximity of the observation radius R to the gravitational radius $R_{\rm h}$. As shown in Melia & Abdelqader (2009), the exact form of $R_{\rm h}$ depends on the constituents of the Universe. For example, in de Sitter space (which contains only a cosmological constant Λ), $R_{\rm h} = c/H_0$, in terms of the (time-invariant) Hubble constant H_0. It is equal to $2ct$ in a radiation-dominated universe and $3ct/2$ when the Universe contains only (visible and dark) matter.

Quite generally, $R_{\rm h}$ is the radius at which a proper sphere encloses sufficient mass-energy to turn it into a Schwarzschild surface for an observer at the origin of the coordinates (see Melia & Shevchuk 2011). That is, R_h is given in all cases by the expression

$$R_{\rm h} = \frac{2GM(R_{\rm h})}{c^2}\ , \tag{4}$$

where $M(R_{\rm h}) = (4\pi/3)R_{\rm h}^3\rho/c^2$, in terms of the total energy density ρ. Thus, $R_{\rm h} = (3c^4/8\pi G\rho)^{1/2}$ which, for a flat universe, may also be written more simply as $R_{\rm h}(t) = c/H_0(t)$.

Although not defined in this fashion, the Hubble radius $c/H_0(t)$ (more commonly encountered when co-moving coordinates are used) is therefore seen to coincide with the gravitational radius $R_{\rm h}$ emerging directly from the Robertson-Walker metric written in terms of R. What this means, of course, is that the Hubble radius is not a mere empirical artifact of the expanding universe, but actually represents the radius at which a sphere centered on the observer contains sufficient mass-energy for its surface to function as a *static* horizon. Of course, that also means that the speed of expansion a distance $R_{\rm h}$ away from us must be equal to c, just as the speed of matter falling into a compact object reaches c at the black hole's horizon (and this is in fact the criterion used to define the Hubble radius in the first place).

The reason why $R_{\rm h}$ is an essential ingredient of the metric written in the form of Equation (2) can be understood in the context of the Birkhoff (1923) theorem and its corollary (see Melia 2007).

[1] It is worth mentioning that prior to the introduction of co-moving coordinates in the 1920's, the cosmological spacetime was actually conventionally expressed in terms of these observer-dependent coordinates (see, e.g., de Sitter 1917.)

The latter states that the metric inside an empty spherical cavity, at the center of a spherically symmetric system, must be equivalent to the flat-space Minkowski metric. Space must be flat in a spherical cavity even if the system is infinite. Thus, the metric between the edge of a cavity and a spherically symmetric mass placed at its center is necessarily of the Schwarzschild type. The worldlines linked to an observer in this region are curved relative to the center of the cavity in a manner determined solely by the mass we have placed there. The implication is that any two points within a medium with non-zero ρ experience a net acceleration (or deceleration) toward (or away) from each other, determined solely by how much mass-energy is contained between them. This is why, of course, the Universe cannot be static, a concept that eluded Einstein himself, since his thinking on this subject preceded that of Birkhoff.

Yet the physical meaning of R_h is still elusive to many, possibly because of the widely held belief that all horizons must necessarily be asymptotic surfaces attained when $t \to \infty$. The so-called event horizon (see Rindler 1956) is indeed of this type, representing the ultimate limit of our observability to the end of time. However, R_h is not in this category—nor should it be. Unlike the event horizon, the gravitational radius is a time-dependent quantity that increases in value at a rate determined by the evolving constituents of the Universe, specifically, the value of the equation-of-state parameter w, defined by the relation $p = w\rho$, where p is the pressure and ρ is the energy density. For some cosmologies, R_h may turn into the event horizon when the cosmic time approaches infinity.

Note from the definition of R_h that $\dot{R}_\mathrm{h}/R_\mathrm{h} = -\dot{\rho}/2\rho$, so with Equation (4)

$$\dot{R}_\mathrm{h} = \frac{3}{2}(1 + w)c \ . \tag{5}$$

Clearly, R_h is constant only for the de Sitter metric, where $w = -1$ and therefore $\dot{R}_\mathrm{h} = 0$. For all other values of $w > -1$, $\dot{R}_\mathrm{h} > 0$. So in ΛCDM, for example, where the Universe is currently dominated by a blend of matter and dark energy, $\dot{R}_\mathrm{h} > 0$. If dark energy were a cosmological constant, however, the Universe would eventually become de Sitter as the density of matter drops to zero, and we would therefore expect R_h to then asymptotically approach a constant value equal to the radius of the event horizon in ΛCDM.

What this means, then, is that the current location of R_h affects what we can observe right now, at time t_0 since the big bang; it is not—and is not meant to be—an indication of how far we will see in our future. One must therefore be careful interpreting spacetime diagrams, such as Figure 3 in van Oirschot et al. (2010). Consider that for an Einstein-de Sitter universe (with $w = 0$, the value used to construct this figure), $\dot{R}_\mathrm{h} = (3/2)c$ so our observations never encounter

R_h because no matter how far light travels, it never overtakes R_h. But that doesn't mean that this must always be the case. In fact, $\dot{R}_h < c$ for any cosmology with $w < -1/3$—which is presumably the situation with ΛCDM. One of the principal goals of this paper is to calculate the actual photon trajectories for well-known FRW cosmologies, such as ΛCDM, in order to demonstrate how and why R_h functions as a limit to our observability today (at time t_0).

3 NULL GEODESICS IN FRW COSMOLOGIES

Let us now derive the equation describing photon trajectories in a cosmology consistent with the FRW metric (Equation 1). From the definition of proper radius, we see that

$$\dot{R} = \dot{a}r + a\dot{r} \ . \tag{6}$$

But the null condition for geodesics (see, e.g., Ellis & Rothman 1993) applied to Equation (1) yields

$$c\,dt = -a(t)\frac{dr}{\sqrt{1-kr^2}} \ , \tag{7}$$

where we have assumed propagation of the photon towards the origin along a radius. The best indications we have today are that the universe is flat so, for simplicity, we will assume $k = 0$ in all the calculations described below, and therefore (for a photon approaching us) $\dot{r} = -c/a$. Thus, we can write Equation (6) as follows:

$$\dot{R}_\gamma = c\left(\frac{R_\gamma}{R_h} - 1\right) \ , \tag{8}$$

where we have added a subscript γ to emphasize the fact that this represents the proper radius of a photon propagating towards us. Note that in this expression, both R_γ and R_h are functions of cosmic time t. The gravitational radius must therefore be calculated according to Equation (5).

The analysis of the WMAP data, within the context of the standard model, points to an age for the Universe of $t_0 \approx 13.7$ billion years. But we also know ρ quite accurately now, and we can use it to calculate $R_h(t_0)$, which appears to be approximately 13.5 billion light-years, equal to ct_0 within the measurement errors. This is the unlikely coincidence we alluded to in the introduction, because there clearly is no particular reason why $\dot{R}_h$ in Equation (5) should be equal to c, especially if w changes with time.

The implausibility of this equality and its possible interpretation have been discussed elsewhere (e.g., Melia & Shevchuk 2011), so we will not dwell on them here. However, we will use the inferred value of R_h today as one of our boundary conditions. In principle, Equation (8) may be integrated either forwards or backwards, but in reality, we are more familiar with the physical

conditions now, at time t_0, so it makes practical sense to begin the solution of this equation at $t = t_0$, where $R_\gamma = 0$ and $R_\mathrm{h}(t_0) = ct_0$.

Not surprisingly, $\dot{R}_\gamma$ equals $-c$ today, since this must represent the actual speed of light measured in our local frame at the origin of our coordinates. Notice, however, that $\dot{R}_\gamma$ differs from $-c$ away from the observer. This type of phenomenon often gives rise to misinterpretation and confusion, stemming from the fact that many consider R to be the actual physical distance measured by a single observer, and that dR/dt is therefore the physical speed. But in general relativity, the proper velocity measured by an individual is actually $v \equiv \sqrt{g_{RR}/g_{TT}}dR/dT$, calculated in terms of the proper distance R and proper time T in his/her frame, where the metric coefficients g_{RR} and g_{TT} are independent of T. Light satisfies the null condition $ds = 0$, and therefore v is *always* equal to c, regardless of whether the frame of reference is inertial or not. Written in terms of the co-moving coordinates, the metric in Equation (1) does not satisfy these conditions, and so $\dot{R}_\gamma$ is generally not equal to c.

This happens because t only represents the physical time on the clock at the observer's location (in other words, $t = T$ only at $R = 0$). If this observer were to measure T at any other position, (s)he would infer a value different from the local proper time t at that radius. And because t is used as a common time everywhere, the quantity $R(t)$ at best represents the sum of all the incremental segments measured by a collaboration of observers—each at time t in their own rest frame—stretched out from one spacetime point to another (see, e.g., Weinberg 1972). Thus, although $R(t)$ is the "proper" distance defined in terms of the cosmic time t, it actually does not represent the physical distance between these two points for an individual observer whose time coordinate is T in his/her frame. A better description for R would be that it represents a "community" distance between two spacetime points and, as a result, $\dot{R}_\gamma$ is not constrained to have the value $-c$ away from the origin.

In the next section, we will discuss the solutions to Equation (8) for various assumed cosmologies, including ΛCDM, the current standard model.

4 RESULTS AND DISCUSSION

The de Sitter universe has no ordinary matter or radiation. Its dynamics is dictated solely by a cosmological constant Λ, which results in an expansion factor $a(t) = e^{H_0 t}$. As we noted earlier (following Equation 3), the gravitational horizon for this spacetime is constant $R_\mathrm{h} = c/H_0$ and $w = -1$. By design, the energy density ρ is also constant. A plot of R_γ, in units of $R_\mathrm{h}(t_0)$, is shown



Figure 1. The null geodesic, R_γ, as a function of cosmic time, t, for de Sitter space. The proper radius is expressed in units of the current gravitational radius $R_{\rm h}(t_0)$, and t is measured as a fraction of the Hubble time t_0.



Figure 2. Same as figure 1, except here for the Einstein-de Sitter universe, consisting entirely of matter.

as a function of t/t_0 in figure 1. In de Sitter, the maximum proper distance of photons reaching us today was $\approx 0.65R_{\rm h}(t_0)$, which occurred at cosmic time $t = 0$. Never along its geodesic path did light's proper distance from us exceed our gravitational horizon today.

This feature is echoed by the Einstein-de Sitter universe, which consists entirely of matter. The cosmological constant and radiation are both absent, so $w = 0$. In this case, the scale factor grows as $a(t) = (3H_0t/2)^{2/3}$ (see, e.g., Melia & Abdelqader 2009), so the density is infinite at $t = 0$ and decreases monotonically as $t \to t_0$. The gravitational radius ($R_{\rm h} = 3ct/2$) grows linearly with time, but is always greater than ct. The null geodesic for this cosmology is shown in figure 2, where we see that R_γ attains its maximum value $\sim 0.3R_{\rm h}(t_0)$ about 2.6 Gyr after the big bang.

When the universal expansion is driven solely by radiation, $w = +1/3$ and $R_{\rm h} = 2ct$. The expansion factor is then given by $a(t) = (2H_0t)^{1/2}$, and the corresponding null geodesic is shown in figure 3. The proper distance R_γ of light attains its maximum value of $\sim 0.4R_{\rm h}(t_0)$ at $t \approx 0.4t_0$.

For a universe dominated by a combination of matter and radiation, the expansion factor also



Figure 3. Same as figure 1, except for a universe dominated by radiation, for which $w = +1/3$ and $R_\mathrm{h} = 2ct$.



Figure 4. Same as figure 1, except for a universe dominated by matter and radiation.

grows monotonically with cosmic time t, though the value of w falls sharply from about $+1/3$ in the early universe (where radiation dominates) and tapers off towards zero as we approach the present time, where the effect of radiation is relatively insignificant. The photon geodesic for this case is shown in figure 4, where we see it attaining its maximum proper distance from us, $\sim 0.3 R_\mathrm{h}(t_0)$, roughly 3 Gyr after the big bang. As with the other cases we've considered, the photon's trajectory never takes it beyond our current gravitational horizon $R_\mathrm{h}(t_0)$.

In ΛCDM, the standard model of cosmology, the density ρ is comprised of three principal components,

$$\rho = \rho_\mathrm{m} + \rho_\mathrm{r} + \rho_\mathrm{de} \tag{9}$$

where, following convention, we designate the matter, radiation, and dark energy densities, respectively, as ρ_m, ρ_r, and ρ_de. We will also assume that these energy densities scale according to $\rho_\mathrm{m} \propto a^{-3}$, $\rho_\mathrm{r} \propto a^{-4}$, and $\rho_\mathrm{de} \propto f(a)$. (If dark energy is indeed a cosmological constant Λ, which we assume for this calculation, then $f(a) =$ constant.) In this case, there is no simple analytical solu-



Figure 5a. The expansion factor $a(t)$ as a function of cosmic time t for the standard model of cosmology, ΛCDM. The constant a_0 represents the value of a at the present time (which in a flat cosmology may also simply be set to 1).



Figure 5b. Same as figure 5a, except here for the value of w time-averaged from 0 to t. Early on, when radiation dominates the equation of state, $\langle w \rangle$ is approximately $+1/3$, but it tapers off and eventually reaches the value $-1/3$ at the present time.

tion for w and $R_{\rm h}(t)$, so these are calculated numerically, along with $R_\gamma(t)$. For completeness, we show the expansion factor $a(t)$ as a function of cosmic time in figure 5a; the value of w, averaged over time from 0 to t, is shown as a function of cosmic time in figure 5b; the gravitational radius $R_{\rm h}$ is shown as a function of t in figure 5c; and the null geodesic itself is plotted in figure 5d. The blending of three different components in ρ produces some discernible differences in the behavior of R_γ compared to the cases we considered previously, but even here, its maximum value was a fraction $\sim$0.45 of today's gravitational radius $R_{\rm h}(t_0)$ at roughly $0.3t_0$ after the big bang.

In all these cases, we have demonstrated through direct computation that null geodesics have never traversed the current gravitational (or Hubble) radius, regardless of what specific cosmology one assumes. In order for light to reach us today, its source radiating at cosmic time t_e must have been located a proper distance $R(t_e)$ away, corresponding to one of the trajectories shown in figures 1,2,3,4, or 5d. In other words, none of the sources whose light we detect today could



Figure 5c. Same as figure 5a, except here for the gravitational radius R_h as a function of t.



Figure 5d. The null geodesic as a function of cosmic time t for ΛCDM.

have been more distant than (in most cases, actually only a fraction ~0.3–0.4 of) our current gravitational horizon $R_\mathrm{h}(t_0)$.

In spite of this, it is sometimes claimed (see, e.g., van Oirschot, Kwan and Lewis 2010) that we see sources from beyond the Hubble radius because its redshift is only ~2, much smaller than sources such as the recombination region that produced the CMB (at a redshift of ~1100 in the standard model). This misconception arises because—written as a proper distance (see Melia & Shevchuk 2011)—R_h is expressed as

$$R_\mathrm{h}(t) = a(t) r_\mathrm{h} \,, \tag{10}$$

where the comoving radius may further be written in the form

$$r_\mathrm{h} = c \int_{t_e}^{t} \frac{dt'}{a(t')} \,. \tag{11}$$

So, for example, if we crudely assume that the Universe's expansion has been driven by a matter-dominated ρ, for which $a(t) = t^{2/3}$, we find that

$$R_\mathrm{h}(t_0) \approx 3ct_0 - 3ct_0{}^{2/3} t_e{}^{1/3} \,, \tag{12}$$

and using the observation that $R_{\rm h}(t_0) \approx ct_0$ (Melia & Shevchuk 2011), with

$$1 + z = \frac{a(t_0)}{a(t_e)} \tag{13}$$

(see, e.g., Weinberg 1972), one easily "finds" a redshift $z_{\rm h}$~1.25 for the Hubble sphere (the value is closer to 2 when using the actual form of $a(t)$ from ΛCDM).

The apparent conflict between this inference and the curves shown in figures 1,2,3,4, and 5d, arises from an incorrect interpretation of $R_{\rm h}(t)$ as a null geodesic, which it is not. In other words, the comoving radius $r_{\rm h}$ in Equation (11), and therefore the proper radius $R_{\rm h}$, do not track the propagation of light through the Hubble flow. Only Equation (8) for $R_\gamma(t)$ can do this because it includes both the effects of an expanding medium and the change in comoving radius r as light approaches us. Instead, $R_{\rm h}(t_0)$ represents today's proper distance to sources that in the past radiated the light we see at t_0 redshifted by an amount $z_{\rm h}$. Therefore, to correctly determine their proper distance at the time the light was produced, one needs to calculate $R_{\rm h}(t_e)$, not $R_{\rm h}(t_0)$, where t_e is the cosmic time corresponding to redshift $z_{\rm h}$. Clearly, $z_{\rm h}$ is that special redshift for which $R_{\rm h}(t_e) = R_\gamma(t_e)$, and there is therefore no conflict between the finite value of $z_{\rm h}$ and the fact that $R_{\rm h}(t_0)$ is the limit to our observability today.

5 CONCLUDING REMARKS

Throughout this paper, we have made a conscious effort to discuss the properties of null geodesics in FRW cosmologies without resorting to conformal diagrams. This approach, also used to great effect by Ellis and Rothman (1993), makes it easier to think in terms of familiar quantities (proper distances and proper time) that are not always straightforward to interpret otherwise. Misconceptions often arise from the misinterpretation of coordinate-dependent effects. In their paper, Ellis and Rothman clearly delineated true horizons from apparent horizons, and extended the definitions, first introduced by Rindler (1956), in a clear and pedagogical manner. In this paper, we have paid particular attention to the gravitational horizon, also manifested as the Hubble radius, which is time-dependent and may or may not turn into an event horizon in the asymptotic future, depending on the equation of state.

It is useful at this point to be absolutely clear about how far light has traveled in reaching us. It is quite evident from our results that sources whose light we see today were at a proper distance $R(t_e) \rightarrow 0$ when $t_e \rightarrow 0$. For the FRW cosmologies we have considered here (which include ΛCDM) the light reaching us today—including that from the recombination region associated with the CMB—has traveled a *net* proper distance of at most ~0.3–0.4ct_0. It is therefore not

correct to claim that the size of the visible universe in these cosmologies is ct_0 (or even greater in some interpretations). Because all causally connected sources in an expanding universe began in a vanishingly small volume as $t \to 0$, the maximum proper distance from which we receive light today must necessarily be less than ct_0, since presumably there were no pre-existing sources at a non-zero proper distance prior to $t = 0$ with which we were in causal contact.

It is true, however, that more and more sources become visible to us as time advances, since for $t > t_0$, the geodesic curves terminating in our future all rise above their current counterparts shown in figures 1,2,3,4, and 5d. In our future, we will see light from sources that radiated at proper distances greater than those shown here. Of course, $R_{\rm h}(t)$ will also continue to increase, and it is not difficult to convince oneself from Equations (5) and (8) that the limits of observability will always be $R_{\rm h}(t)$, since $R_\gamma(t_e)/R_{\rm h}(t)$ is smaller than 1 for all $t_e < t$. To see this, let us first find the emission time $t_{e,\,\rm max}$ at which $R_\gamma(t_e)$ attains its maximum value. Clearly, this happens when $\dot{R}_\gamma = 0$, which means that $R_\gamma(t_{e,\,\rm max}) = R_{\rm h}(t_{e,\,\rm max})$. But for all $w > -1$, Equation (5) shows that $\dot{R}_{\rm h} > 0$, and therefore $R_{\rm h}(t) > R_{\rm h}(t_{e,\,\rm max})$ for $t > t_{e,\,\rm max}$, which also means that $R_{\rm h}(t) > R_\gamma(t_{e,\,\rm max})$ for all $t > t_{e,\,\rm max}$.

ACKNOWLEDGMENTS

This research was partially supported by ONR grant N00014-09-C-0032 and an NSF Graduate Fellowship at the University of Arizona, and by a Miegunyah Fellowship at the University of Melbourne. We are particularly grateful to Amherst College for its support through a John Woodruff Simpson Lectureship. Finally, we wish to thank Roy Kerr for many helpful and enjoyable discussions.

REFERENCES

Birkhoff, G. 1923, Relativity and Modern Physics, Harvard University Press

Brax, P., van de Bruck, C., Davis, A.-C., Khoury, J., & Weltman, A. 2004, Phys. Rev. D., 70, 123518

Capozziello, S., Cardone, V. F., Carloni, S. & Troisi, A. 2003, IJMP-D, 12, 1969.

Carroll, S. M., Duvvuri, V., Trodden, M. & Turner, M. S. 2004, Phys. Rev. D, 70, 043528

Davis, T. M. & Lineweaver, C. H. 2004, PASA, 21, 97

de Sitter, W. 1917, Proc. Akad. Weteusch. Amsterdam, 19, 1217

Ellis, G.F.R. & Rothman, T. 1993, Am. J. Phys. 61, 883

Khoury, J. & Weltman, A. 2004, Phys. Rev. Lett., 93, 171104

Lima, J. A. S. 2007, arXiv:0708.3414 (unpublished)

Melia, F. 2003, The Edge of Infinity: Supermassive Black Holes in the Universe, Cambridge University Press, pp. 119–128

Melia, F. 2007, MNRAS, 382, 1917

Melia, F. 2009, Int. J. Mod. Phys. D, 18, 1113

Melia, F. & Abdelqader, M. 2009, Int. J. Mod. Phys. D, 18, 1889

Melia, F. & Shevchuk, A.S.H. 2011, MNRAS, in press

Mota, D. F. & Barrow, J. D. 2004, MNRAS, 349, 291

Nojiri, S. & Odintsov, S. D. 2003, Phys. Rev. D, 68, 123512

Ratra, B. & Peebles, J. 1988, Phys. Rev. D, 37, 3406

Rindler, W. 1956, MNRAS, 116, 662

Spergel, D. N. et al. 2003, ApJS, 148, 175

Tegmark, M., et al. 2004, Phys. Rev. D, 69, 103501

van Oirschot, P., Kwan, J. & Lewis, G. F. 2010, MNRAS, in press (arXiv:1001.4795v1)

Weinberg, S. 1972, Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity, (Wiley, New York)

Wetterich, C. 1988, Nucl. Phys. B, 302, 668